

「香港華人婦女與性」

(育齡期及更年期婦女)

研究結果發佈

Findings of Studies on

Sexuality of Hong Kong Chinese Women (Reproductive Age and around Menopause)



Maturitas 74 (2013) 190-195

ORIGINAL RESEARCH—WOMEN'S SEXUAL HEALTH

Sexual Behavior and Symptoms among Reproductive Age Chinese Women in Hong Kong

Sue Seen-Tsing Lo, MD and Wai-Ming Kok, MSc

The Family Planning Association of Hong Kong, Hong Kong SAR, China

DOI: 10.1111/jsm.12508

ABSTRACT-

Introduction. As sexual medicine evolves, much advancement has been achieved in understanding male sexuality and treating male sexual dysfunction. Less is known about female sexual pattern, the prevalence of sexual problems, and their correlation with confounding factors.

Aim. To enhance our understanding of female sexuality and the risk factors that contributed to sexual problems in reproductive age women.

Method. A cross-sectional survey was conducted in family planning and prepregnancy checkup clinics from December 2007 to December 2009, with 2,146 sexually active Chinese women aged 21 to 40 years completed the entire questionnaire.

Main Outcome Measures. Prevalence of sexual symptoms, coital frequency, and other sexual behavior-related activities were measured.

Results. Overall, 59.0% of respondents had at least one sexual problem. In this sample, 31.8% of respondents reported no desire; 31.7% had arousal problems; 40% had anorgasmia, and 33.8% experienced coital pain for at least 3 months within the past 1 year. Chi-square test showed significant correlation among the four types of sexual problems(P < 0.001). Univariate regression model showed that all sexual symptoms were significantly correlated with unidirectional coitus initiation, low coital frequency, and low foreplay enjoyment. Loglinear model revealed that desire, arousal, and orgasmic problems were correlated with low foreplay enjoyment. Arousal problem was correlated with high acceptance toward pornography and history of medical disease. Coital pain was correlated with secondary education and planning to have more children. Both unidirectional coitus initiation and low coital frequency were major contributors to all four sexual symptoms.

Conclusions. Sexual problem is a prevalent health issue among reproductive age women. A number of risk factors are identified, which provide useful direction to the design of counseling and education materials that might help to enhance sexual performance in women. Lo SST and Kok WM. Sexual behavior and symptoms among reproductive age Chinese women in Hong Kong. J Sex Med 2014;11:1749–1756.

Key Words. Female Sexuality; Female Sexual Dysfunction; Risk Factors; Sexual Activities; Epidemiology



Contents lists available at SciVerse ScienceDirect

Maturitas

journal homepage: www.elsevier.com/locate/maturitas



Sexuality of Chinese women around menopause

Sue Seen-Tsing Lo*, Wai-Ming Kok

The Family Planning Association of Hong Kong, G/F, 8/F, 9/F, 10/F Southorn Centre, 130 Hennessy Road, Wanchal, Hong Kong Special Administrative Region

ARTICLE INFO

Article history; Received 28 May 2012 Received in revised form 31 October 2012 Accepted 16 November 2012

Keywords; Female sexual dysfunction Menopause Sexual satisfaction Sexual attitude Sexual activities

ABSTRACT

Objective: To study the sexual activities and prevalence of sexual dysfunctions in midlife Chinese women and their correlations with demographic factors, sexual dissatisfaction and interpersonal difficulty. Study design: This is a cross-sectional survey of a convenience sample of women aged 40-60, who requested gynecological checkup or attend social activities at Women's Club.

Main outcome measures: Sexual activities, sexual dysfunctions, sexual dissatisfaction, demographic factors and interpersonal difficulty were assessed by self-administered questionnaire.

Results: Among 371 eligible subjects, 22.4% and 39.6% women had low intimacy and coitus frequency (0 to <12 acts in one year), respectively. The odds ratios for low coital frequency in the natural menopausal and surgical menopausal subgroups were 3.00 and 5.09, respectively (95% confidence interval: 1.73–5.19 and 1.77–14.69, respectively). Overall, 77.2% women had at least one type of sexual dysfunctions; this proportion was highest in the surgically menopausal subgroup (88.9%) followed by the naturally menopausal subgroup (78.2%) and the premenopausal subgroup (72.2%) (p=0.003). No lubrication (42.9%) was the commonest sexual dysfunction and predominantly affected naturally and surgically menopaused women (p=0.001). Sexual dysfunction was the major contributor to sexual dissatisfaction (0.80), followed by interpersonal difficulty (0.2). Arousal disorder was the pivot of interaction between sexual dissatisfaction, menopausal status and low coital frequency. Conclusions: Chinese women had fewer intimate contacts and less coitus when menopause progressed. No lubrication was the commonest sexual dysfunction and predominantly affected menopaused women. Our model showed that sexual dysfunction is the main contributor to sexual dissatisfaction.

© 2012 Elsevier Ireland Ltd. All rights reserved.

研究方法

2007年12月至2009年12月 不記名問卷

21-40歲 節育指導服務 懷孕前準備服務 40-60歲 女性保健服務 婦女會



「香港華人婦女與性」 (育齡期及更年期婦女) 研究結果發佈

Findings of Studies on
Sexuality of Hong Kong Chinese Women
(at Reproductive Age and around Menopause)



性生活模式

每年至少12次	21-40歲 (2146人)	40-60歲 (371 人)
性幻想	77.7%	35.9%
與伴侶有親密接觸	93.0%	77.9%
性交	81.9%	60.4%





「香港華人婦女與性」 (育齡期及更年期婦女) 研究結果發佈

Findings of Studies on Sexuality of Hong Kong Chinese Women (at Reproductive Age and around Menopause)



性生活模式

毎年至少12次	未收經 (117人)	收經過渡期 (125人)	已收經 (111人)	手術絕經 (18人)
性幻想	35.1%	43.2%	28.8%	33.3%
與伴侶有親密 接觸	82.9%	76.5%	66.9%	52.6%
性交	71.8%	66.4%	46.0%	(33.3%)



「香港華人婦女與性」 (育齡期及更年期婦女) 研究結果發佈

Findings of Studies on Sexuality of Hong Kong Chinese Women (at Reproductive Age and around Menopause)







在性交時遇到的問題

過去一年,至少連續三個月都有 問題	21-40歲
欠缺性幻想	31.8%
不感興奮	31.7%
無高潮	40.0%
疼痛	33.8%
至少有一個問題	





「香港華人婦女與性」 (育齡期及更年期婦女) 研究結果發佈

Findings of Studies on Sexuality of Hong Kong Chinese Women (at Reproductive Age and around Menopause)



過去一年,至少連續 三個月都有問題	40-60歲	未收經/過渡中/已收經/手術
欠缺性幻想	35.6%	32.5 / 36.0 / 37.8 / 38.9%
不感興奮	38.3%	33.3 / 38.4 / 43.2 / 38.9%
陰道乾涸	42.9%	29.1 / 44 / 55 / 50%
無高潮	41.0%	35.9 / 41.6 / 45 / 44.4%
疼痛	31.5%	25.7 / 30.4 / 38.7 / 33.4%
至少有一個問題 -	77.2%	72.2 / 78.2 / 79.3 / 88.9%





「香港華人婦女與性」 (育齡期及更年期婦女) 研究結果發佈

Findings of Studies on
Sexuality of Hong Kong Chinese Women
(at Reproductive Age and around Menopause)



性功能障礙的誘因

21-40歲	40-60歲
單(男)方主導性交要求	更年期
很少有性交	
不享受前戲	





「香港華人婦女與性」 (育齡期及更年期婦女) 研究結果發佈

Findings of Studies on Sexuality of Hong Kong Chinese Women (at Reproductive Age and around Menopause)



性功能障礙的影響

21-40歲	40-60歲
低生育率	性生活不滿意
降低女性的自主性	萎縮性陰道炎
影響感情、婚姻	





「香港華人婦女與性」 (育齡期及更年期婦女) 研究結果發佈

Findings of Studies on Sexuality of Hong Kong Chinese Women (at Reproductive Age and around Menopause)



消除障礙

21-40歲	40-60歲
增強夫婦間的性溝通	保持性生活
克服談性的尷尬	至少每星期一次
輪流採取主動	潤滑劑、荷爾蒙
注重、享受前戲	了解性反應隨年齡的變化
增加情趣	多溝通 、互相體諒





「香港華人婦女與性」 (育齡期及更年期婦女) 研究結果發佈

Findings of Studies on Sexuality of Hong Kong Chinese Women (at Reproductive Age and around Menopause)





香港家庭計劃指導會 The Family Planning Association of Hong Kong

性生活

講座系列



2014年7月至12月

家計會性教育網頁 www.famplan.org.hk/sexedu

課程查詢及報名 家計會教育組 29197734

課程查詢及報名 家計會教育組 29197734

全人《性生活講座系列》簡介

我們相信,和諧協調的性關係,可以帶來身心的愉悦和滿足。雖說性 是人的本能,但美滿的性生活仍需要伴侶間多方面的配合,這包括對性有 正確的認識,抱開明坦誠的性態度,掌握得宜的性技巧,並要懂得互相溝 通和配合,才不會因一些誤解、甚至無知而產生焦慮,及性生活不協調。

性生活講座系列,分基礎知識篇及親密情趣篇,旨在增強參加者對性 生活的基本認識,希望可幫助參加者順利展開性生活,並提高其性生活質 素,增添情趣。

基礎知識篇

較適合準備或 剛展開性生活的人士

內容包括 認識男女生殖系統 男女性反應差異 性交過程及基本姿勢 性交協調技巧 如何選擇避孕方法 和諧性生活要素

親密情趣篇

較適合想改善性生活,或 在性生活方面出現疑難的人士

內容包括

如何促進性生活的溝通 提升性刺激的技巧 如何增添性情趣 懷孕期及分娩後的性生活 性功能障礙

主講者皆為資深性教育工作者,任職香港家計會多年,經常主持各類 型性教育講座及培訓活動,發展性教育資源,亦常於傳媒及網上解答公眾 人士性問題,並為本地報章雜誌撰寫性教育文章,著作有〈性生活、情趣 小百科>、<男性健康>、<識婚男女>等。

凡成功参加講座者,可獲騰本會出版之〈性生活・情趨小百科〉或〈議婚男女 >或《婚前及懷孕前的準備DVD>,以及本會活動及資源館個人會籍一年。



「香港華人婦女與性」

(育齡期及更年期婦女)

研究結果發佈

Findings of Studies on Sexuality of Hong Kong Chinese Women (at Reproductive Age and around Menopause)









1999

Extending Your Golden Years:

A Healthy & Happy Menopause







「香港華人婦女與性」 (育齡期及更年期婦女) 研究結果發佈

Findings of Studies on Sexuality of Hong Kong Chinese Women (at Reproductive Age and around Menopause)

24 • 9 • 2014



2001





個人資料·絕對保密 收費如有更改,想不另行通知

總會辦事處:香港灣仔軒尼詩道130號修頓中心十樓

網址:www.famplan.org.hk 電子郵箱:fpahk@famplan.org.hk





看港家原計劃推導會是一所非牟利的志願機構,我們需要社會各界熱心 人士、惠書團體等的支持和歐低預助,使服務發展更讓完善。我們亦敬認 例加入股為裁工。詳問請考覽本會調養、www.famplan.org/h·妻於電2572-2272卷編。

我們希望本會資料單級能夠符合你的需要·如有任何改善建議·歌迎致電 2575-4477和我們聽信·

持有社會福利署或關前管理局發出「醫療費用減免證明書」的人士,可於整 記削出示有效文件,申請部免部仍服務收費。

© 2014 香港家庭計劃指導會 歐遊復製此單張內容 作非牟利教育用途

敬請留意本會網站最新資料





2007

「性」是與生俱來的,圓滿的性生活是維繫美滿婚姻生活的重要元素。家計會「性治療」服務透過專業的醫療、輔導及啟導服務,協助夫婦克服性障礙,重拾性愛的親密和歡愉,並建立快樂、健康、和諧的美滿婚姻生活。




「香港華人婦女與性」

(育齡期及更年期婦女) 研究結果發佈

Findings of Studies on Sexuality of Hong Kong Chinese Women (at Reproductive Age and around Menopause)















「香港華人婦女與性」 (育齡期及更年期婦女) 研究結果發佈

Findings of Studies on Sexuality of Hong Kong Chinese Women (at Reproductive Age and around Menopause)

